



## GENOTYPIC RESPONSE OF SORGHUM (*SORGHUM BICOLOR*) MUTANT LINES FOR YIELD AND QUALITY OF FODDER UNDER SUB-HUMID SOUTHERN PLAIN

JOGENDRA SINGH, LATA CHAUDHARY AND NEELAM SHEKHAWAT

Department of Plant Breeding and Genetics, Rajasthan College of Agriculture, Maharana Pratap University of Agriculture and Technology, Udaipur (Rajasthan)-313 001, India

Received: 30.01.2014

Revised accepted: 28.02.2014

### ABSTRACT

#### Keywords:

Multicut forage sorghum,  
Mutant lines,  
SSG 59-3.

An investigation was carried out with multicut forage sorghum variety SSG 59-3 and its 15 mutants derived from gamma irradiation to identify the superior mutant genotypes for high fodder yield and quality in sorghum. Differences among the genotypes were found significant for all the quality traits and most of the yield traits studied at different cut(s). The mutant genotypes SSG 226 was the best performer for both quality and fodder yield and, another two mutant genotypes SSG 231 and SSG 222 was also good for fodder yield and quality, respectively but it perform poorer for vice-versa. The genotype SSG 226 produced green fodder yield (3.33, 1.52 and 0.95 g/plant/day) and dry fodder yield (1.03, 0.61 and 0.42 g/plant/day) at first, second and third cuts, respectively with crude protein (8.18) along with desirable lowest crude fibre (30.60) and highest ash content (8.39) at first cut. Taking a better variety in respect to fodder yield and nutritional contents, the genotype SSG 226 should be preferred over the tested mutant genotypes for forage purpose.

### INTRODUCTION

Availability of adequate quantity of quality feed and fodder for livestock is essential for sustaining the livestock productivity. Due to increasing pressure on land for growing food grains, oil seeds, and pulses, fodder production generally gets lower priority. With about 2.29% share of the land area of the world, India is maintaining about 10.71% world's livestock (SIA, 2013). Further, inadequate production and availability of improved fodder seeds, diverse uses of agriculture crop residues (paper industry, packaging, etc.), area has been declined under coarse cereals which are also used as feed for last 30 years, a substantial amount of crop residues is burnt by the farmers after harvesting the main crops like wheat and paddy, subsequently, the gap between the demand and supply of fodder is increasing. Fodder and feeds are the major inputs in animal production especially in milch animals, which account for about 60-70% of total cost of milk production. The present availability of green fodder is about 513 million tonnes projecting a deficit of 53% and that of dry fodder is around 400 million tonnes against the requirement of 676 million tonnes (Mukherjee *et al.*, 1998). At present, fodder is being cultivated only on 4% of gross cropped area, which is not adequate to meet the

requirement of the livestock (State of Indian Agriculture, 2012-13).

The forage crops are the cheapest source of animals feed and therefore, taken as foundation of livestock industry. The demand for livestock products is continuously rising due to their regular use in human diets. It has been estimated that need for forage crops up to 2050 will increase two to three folds in Asian countries (Devendra and Leng, 2011). To overcome such situation, genetically stable genotypes having good nutritional value and high fodder yield potential are urgently needed. Sorghum fodder plays an important role in the health and nutrition of the large population of livestock in the country by providing nutritive fodder. Sorghum is an important crop widely grown for grain and fodder with a greater emphasis on fodder particularly in semi-arid tracts. Sorghum produces a tonnage of dry matter having proportions of digestible nutrients (50%), crude protein (8%), fat (2.5%) and nitrogen free extracts (45%) (Azam *et al.*, 2010). It can be used fresh as well as stored in form of silage and hay for future use. As a result of crop improvement programme, a number of promising strains of plants with diversified morphological and quality traits are available for general cultivation (Hussain *et al.*, 1995).

\*Corresponding author email: jschouhanpbg@gmail.com

The changes in genetic material of crops resulted wide variations in the morphological and forage quality traits (Alias *et al.*, 2010; Ullah *et al.*, 2007). Therefore, genetic improvement of crop is basically aimed to enable the crop to survive in environmental vagaries. The planned study was conducted with the objective to understand the response of mutant lines under the Udaipur conditions and identification of suitable mutant line for growing under such conditions and even some of the mutant lines may be used in the breeding programme for some specific traits for further development of suitable genotypes for high quality forage production.

#### MATERIALS AND METHODS

An experiment on forage sorghum [*Sorghum bicolor* (L.) Moench] was conducted during summer-2010 at Instructional Farm of Rajasthan College of Agriculture, MPUAT, Udaipur (Rajasthan). Udaipur is situated at South-Eastern part of Rajasthan at an altitude of 579.5 m MSL and at 24° 35' N latitude and 74° 42'E longitude. The region falls under agro-climatic zone IV a

(Sub- humid Southern Plain and Aravalli Hills) of Rajasthan. The experiment was conducted during 24<sup>th</sup> May to 21<sup>st</sup> October, 2010 on clay loam soil under irrigated conditions. The mean meteorological data pertaining to crop season are given in Table-1. The experimental material comprised of 15 mutant lines in M<sub>5</sub> generation, viz., SSG 222, SSG 224, SSG 225, SSG 226, SSG 227, SSG 231, SSG 232, SSG 233, SSG 234, SSG 236, SSG 241, SSG 244, SSG 253, SSG 256 and SSG 263 obtained through the use of gamma-rays, along with its parent SSG 59-3 (a popular variety of multicut forage sorghum), were planted in randomized block design with three replications. Each genotype had four rows of 4 m length with 25 cm row to row and 15 cm plant to plant spacing. The recommended cultural practices were adopted for raising the good crop. The observations were recorded for 14 different characters at different cut (s) on five randomly selected plants for each genotype in each replication.

**Table 1 Week-wise average of meteorological data during the crop growth season (Kharif-2010)**

Standard Week No.	Dates	Temperature (°C)		Relative Humidity (%)		Sunshine (hrs)	Rainfall of week (mm)
		Max.	Min.	Max.	Min.		
20	14 May - 20 May	42.2	27.7	32	16	9.9	0.0
21	21 May - 27 May	43.1	28.8	36	22	9.1	0.0
22	28 May - 3 June	39.6	28.2	51	25	8.6	0.0
23	4 June - 10 June	35.5	25.2	65	36	8.3	15.6
24	11 June - 17 June	39.8	28.0	48	31	6.5	0.0
25	18 June - 24 June	39.6	28.1	52	31	8.0	0.0
26	25 June - 1 July	37.4	25.6	75	38	8.8	19.5
27	2 July - 8 July	33.3	24.1	84	65	5.0	120.0
28	9 July - 15 July	32.1	25.4	80	62	4.7	0.0
29	16 July - 22 July	33.0	24.9	88	71	4.0	87.2
30	23 July - 29 July	28.7	23.5	93	84	2.5	113.9
31	30 July - 5 Aug.	30.3	23.4	91.4	74.7	3.3	154.8
32	6 Aug. - 12 Aug.	29.5	23.9	92.3	83.0	3.1	69.2
33	13 Aug. - 19 Aug.	30.8	24.0	92.7	75.9	3.2	16.5
34	20 Aug. - 26 Aug.	30.8	23.6	86.0	66.1	5.9	27.2
35	27 Aug. - 2 Sept.	31.0	23.3	92.6	79.3	3.9	55.7
36	3 Sept. - 9 Sept.	30.4	23.2	93.4	81.7	3.6	100.3
37	10 Sept. - 16 Sept.	28.8	22.8	91.6	77.6	2.5	13.2
38	17 Sept. - 23 Sept.	32.1	18.5	81.9	39.4	8.1	0.0
39	24 Sept. - 30 Sept.	32.8	18.4	77.6	32.4	8.8	0.0
40	1 Oct. - 7 Oct.	34.5	18.3	76.1	29.3	9.2	0.0
41	8 Oct. - 14 Oct.	33.9	17.7	77.1	31.0	8.3	0.0

Source: Agro-meteorological Observatory, Instructional farm, RCA, Udaipur

Observations on green fodder yield and related components were recorded at 60 days after sowing (DAS) during first cut, 45 days after first cut (DAFC) during second cut and 45 days after second cut (DASC) during third cut while number of tillers per plant was recorded at cutting stage of only second and third cuts. Quality parameters viz., crude protein, crude fibre, ether extract, nitrogen free extract, ash and TDN were estimated from dry fodder at first cut only, while N content in plants were estimated from dry fodder from all the three cuts. Besides, HCN content in plants was calculated at 30 DAS, 30 DAFC and 30 DASC, respectively using Picric acid method given by Hogg and Ahlgren (1942). Fresh plant samples of the various genotypes were collected from each replication and evaluated for the nitrogen content and remaining forage quality parameters (crude protein, crude fibre, nitrogen free extracts, ether extract and total ash). Plant samples were chopped mixed thoroughly and grind to fine powder and were divided into three groups for estimations of the following quality components by using proximate analysis (AOAC, 1975). The data recorded were subjected to analysis of variance (Steel *et al.*, 1997) for the mentioned characteristics to determine the significance of differences among genotypes.

## RESULTS AND DISCUSSION

The analysis of variance revealed significant differences among the genotypes for all the quality traits and most of the morphological and yield traits studied at different cut(s) Table 1.

### Growth parameters

As shown in Table-1, the difference among the genotypes with respect to the plant height was significant at third cut only while it was non-significant at first and second cut. The average plant height was 244.83 cm at first cut, 140.17 cm at second cut and 85.77 cm at third cut. At first cut, SSG 241 (271.33) had maximum plant height values followed by SSG 233 (256.00), SSG 263 (256.00) and SSG 225 (255.00) while lowest values was observed in SSG 256 (222.00). At second cut, maximum plant height was observed in SSG 224 (176.33) followed by SSG 226 (167.67) and SSG 232 (160.00) while it was minimum for SSG 263 (90.67). At third cut, maximum plant height was observed in check SSG 59-3 (116.67) followed by SSG 226 (115) and SSG 253 (113.33) while, genotype SSG 233 showed minimum plant height (47.67 cm). It may be due to varietal effect with its morphological influenced (Ayub *et al.*, 2010; Al-Din *et al.*, 2012; Ayub *et al.*, 2012; Singh *et al.*, 2013).

As shown in Table-1, the difference among the genotypes with respect to the number of leaves per plant was significant at first cut only while it was non-significant at second and third cut. The average number of leaves was 11.52 at first cut, 9.35 at second cut and 6.96 at third cut. Data from first cut revealed that maximum number of leaves was observed in SSG 222

(12.80) followed by SSG 244 (12.20) while SSG 263 was equal to check SSG 59-3 (12.13) for number of leaves and its value was minimum in SSG 224 (10.53). At second cut, values of number of leaves varied from 8.33 (SSG 233) to 10.33 (SSG 226) and SSG 232 (10.20) also possessed higher number of leaves than check SSG 59-3 (10.00). At third cut, highest number of leaves was exhibited by check SSG 59-3 (8.33) followed by SSG 226 (8.00) and lowest value showed in SSG 234 (5.67). The significant differences among sorghum cultivars has also been previously reported by Nabi *et al.* (2006), Ayub *et al.* (2012), Seetharam and Ganesamurthy (2013) and Singh *et al.* (2013) which probably due to genetic make of genotypes under investigation.

As shown in Table 1, the difference among the genotypes with respect to the stem girth was significant at third cut only while it was non-significant at first and second cut. The average values of stem girth were 4.13 cm at first cut, 3.56 cm at second cut and 2.60 cm at third cut. Highest stem girth values were obtained from the SSG 231 (4.67) in first cut while SSG 222 (4.53) and SSG 253 (4.40) also possessed higher value of stem girth than check SSG 59-3 (4.37). Thinnest stem was observed in SSG 224 (3.67). At second cut, highest stem girth values were obtained from the SSG 231 (4.00) followed by SSG 232 (3.93) and SSG 263 (3.80). Thinnest stem was observed in SSG 222 (3.10). At third cut, highest stem girth values were obtained from the SSG 253 (3.47) followed by SSG 244 (3.0), SSG 227 (2.95) and SSG 232 (2.93). Thinnest stem was observed in SSG 234 (2.07). Our results for stem diameter has also been confirmed by the findings of Ayub *et al.* (2012); Ghasemi *et al.* (2012) where a range of stem diameter was observed for sorghum cultivars.

As shown in Table1, the differences among the genotypes with respect to the number of tillers per plant were significant at both second and third cut. The average of number of tillers per plant was 1.64 at second cut and 2.62 cm at third cut. Maximum number of tillers produced by check SSG 59-3 (2.55 and 3.89) followed by SSG 226 (2.44 and 3.66) at second and third cut, respectively. Lowest tillers number was found in SSG 244 at both the cuts. Our result has also been confirmed by the findings of Ghasemi *et al.* (2012) and Singh *et al.* (2013) for tiller number in sorghum.

As shown in Table-1, the difference among the genotypes with respect to the green fodder yield/plant/day was significant at third cut only while it was non-significant at first and second cut. The average values of green fodder yield/plant/day were 3.02 g at first cut, 1.32 g at second cut and 0.60 g at third cut. At first cut, highest green fodder yield value was produced by SSG 231 (3.99) followed by SSG 263 (3.88), SSG 253 (3.59) and SSG 244 (3.55) while low green fodder yield/plant/day was produced by SSG 241, SSG 233, SSG 222 and SSG 225 than check. At second cut,

maximum green fodder yield/plant/day value was observed in SSG 244 (1.64) while minimum value showed by SSG 232 (1.02). Besides, two other genotypes SSG 256 (1.63) and SSG 226 (1.52) also yielded higher than check SSG 59-3 (1.41). At third cut, green fodder yield varied from 0.30 (SSG 263) to 0.95 g (SSG 226). Besides, three other genotypes SSG 241 (0.90), SSG 224 and SSG 253 (0.89) also showed higher yield than check SSG 59-3 (0.81). The significant differences in green forage yield among sorghum cultivars have also been undertaken by Chughtai *et al.* (2007); Ayub *et al.* (2012); Ghasemi *et al.* (2012); Singh *et al.* (2013).

#### **Dry fodder yield/day/plant (g)**

As shown in Table-1, the difference among the genotypes with respect to the dry fodder yield/plant/day was non-significant at all the three cuts. The average values of dry fodder yield/plant/day were 0.74 g at first cut, 0.51 g at second cut and 0.33 g at third cut. At first cut, highest dry fodder yield/plant/day value was produced by SSG 263 (1.24) followed by SSG 253 (1.12), SSG 234 (1.04), SSG 231 and SSG 226 (1.03) while low green fodder yield/plant/day was produced by SSG 222, SSG 233 and SSG 241 than check. At second cut, highest and lowest dry fodder yield was recorded in SSG 225 (0.68) and SSG 233 (0.38), respectively. Besides, 4 other genotypes SSG 226 (0.61), SSG 256 (0.59) SSG 231 and SSG 244 (0.53) were also yielded higher than check SSG 59-3 (0.51). At third cut, data revealed that maximum dry fodder yield was recorded in SSG 224 (0.49) whereas minimum was in SSG 234 and SSG 263 (0.15). Besides, 3 other genotypes SSG 226 (0.42), SSG 225 (0.37) and SSG 253 (0.36) also exhibited higher yield than check SSG 59-3 (0.33). Significant variations among sorghum genotypes for dry matter production have already been reported in studies conducted by Yousef *et al.* (2009); Ayub *et al.* (2012); Ghasemi *et al.* (2012); Singh *et al.* (2013).

#### **Quality parameters**

As shown in Table-1, the differences among the genotypes with respect to all quality parameters were significant at different cut (s) as these were studied.

#### **Nitrogen content in plant (%)**

As shown in Table-3, the difference among the genotypes with respect to the nitrogen content in dry fodder was significant at all the three cuts. The average values of nitrogen content in plant were 1.14 at first cut, 0.95 at second cut and 0.77 at third cut. At first cut, the maximum nitrogen content in plant was exhibited by check SSG 59-3 (1.41) followed by SSG 222 (1.35), SSG 226 (1.31) and SSG 233 (1.28) while minimum value was observed in SSG 263 (0.89). Data from second cut revealed that the maximum nitrogen content in plant was observed in check SSG 59-3 (1.20) followed by SSG 222 and SSG 226 (1.14) while minimum value showed in SSG 263 (0.75). At third cut, the maximum nitrogen content was exhibited in check

SSG 59-3 and SSG 226 (0.96) followed by SSG 222 (0.92) and SSG 233 (0.88) while minimum N content was recorded in SSG 232 and SSG 263 (0.62)..

#### **Crude protein (%)**

Data from first cut (Table-3) revealed that average value of crude protein in dry fodder was 7.14 % and check SSG 59-3 exhibited maximum crude protein (8.79) followed by SSG 222 (8.45), SSG 226 (8.18) and SSG 233 (8.03) while minimum value was observed in SSG 263 (5.54). The higher protein contents in dry matter ultimately will result higher protein yield on unit area. The significant differences in crude protein contents in dry matter of various genotypes have also been confirmed by Nabi *et al.* (2006); Tauqir *et al.* (2009); Ayub *et al.* (2012); Bibi *et al.* (2012). The difference among genotype may be due to relative contribution of leaves to total biomass and concentration of protein in dry matter.

#### **Crude fibre (%)**

Data from first cut (Table-3) revealed that the average value of crude fibre content in dry fodder was 31.89%. Desirable lower value was observed in SSG 226 (30.60) followed by check SSG 59-3 (30.65) and SSG 233 (30.78) while, maximum values were observed in SSG 263 (33.50). Significant differences among sorghum genotypes for crude fibre have already been confirmed by studies conducted by Nabi *et al.* (2006); Bibi *et al.* (2012).

#### **Ether extract (%)**

Data from first cut (Table-3) revealed that the average value of ether extract in dry fodder was 1.74 %. Maximum ether extract was calculated in check SSG 59-3 (1.85) followed by SSG 222 (1.82), SSG 244 (1.81), SSG 226 and SSG 233 (1.80) while, minimum value was observed in SSG 263 (1.65). Similar results were also reported by Bibi *et al.* (2012); Singh *et al.* (2013).

#### **Ash content (%)**

Data from first cut (Table-3) revealed that the average value of ash content in dry fodder was 7.63%. Maximum ash content was observed in SSG 226 (8.39) followed by check SSG 59-3 (8.33), SSG 222 (8.20) and SSG 244 (8.15). Significant variations in ash contents among tested genotypes suggested differences in nutrient absorption from soil and utilization within the plants. The results are consistent with those of Nabi *et al.* (2006); Bibi *et al.* (2012).

#### **Nitrogen free extract (%)**

Data from first cut (Table-3) revealed that the average value of nitrogen free extract in dry fodder was 51.56%. Desirable minimum value of nitrogen free extract was observed in SSG 222 (50.30) followed by check SSG 59-3 (50.40) and SSG 244 (50.79). Maximum values were observed in SSG 263 (52.39). Similar results also reported by Bibi *et al.* (2012); Singh *et al.* (2013) in sorghum.

**Table 2 Mean square for various characters at different cut (s) in forage sorghum**

Characters	Replication	Genotype	Error
	[2]	[15]	[30]
Plant height - I cut	1570.3959*	474.6667	436.5
Plant height - II cut	193.0833	1484.4889	1060.0
Plant height - III Cut	4265.8960**	1491.7208**	341.4
Number of leaves/plant - I cut	12.0808**	1.1387**	0.4062
Number of leaves/plant - II cut	4.4033**	0.9026	0.6158
Number of leaves/plant - III Cut	5.1458*	1.6389	1.101
Stem girth - I cut	0.6175*	0.2371	0.1244
Stem girth - II cut	0.4075	0.1661	0.1337
Stem girth - III Cut	0.7656*	0.4624**	0.1554
Number of tillers/plant - II cut	0.0281	0.4843**	0.05482
Number of tillers/plant - III Cut	0.1339	1.1858**	0.2116
Green fodder yield/plant/day - I cut	2.6612*	0.9885	0.6823
Green fodder yield/plant/day - II cut	1.8404**	0.1022	0.169
Green fodder yield/plant/day - III Cut	0.1601*	0.1543**	0.03559
Dry fodder yield/plant/day - I cut	0.2721*	0.0890	0.08117
Dry fodder yield/plant/day - II cut	0.7058	0.2712	0.2386
Dry fodder yield/plant/day - III Cut	0.4130	0.2998	0.2547
N content in plant - I cut	0.0047	0.0639**	0.001435
N content in plant - II cut	0.0009	0.0519**	0.0007865
N content in plant - III cut	0.0010	0.0400**	0.0007954
Crude protein - I cut	0.1600	2.4789**	0.05982
Crude fibre - I cut	0.0878	2.2437**	0.2245
Ether extract - I cut	0.0008	0.0116**	0.0003463
Ash - I cut	0.1739	0.6691**	0.09449
Nitrogen free extract - I cut	1.0754**	1.3782**	0.1211
TDN - I cut	0.0174	0.4607**	0.005935
HCN content - I cut	1.5625	4553.2984**	122.7
HCN content - II cut	16.1458	3653.0208**	71.15
HCN content - III cut	118.7500	865.4167**	52.08

\*, \*\* Significant at 5 % and 1 % level, respectively.

**TDN (%)**

Data from first cut (Table-3) revealed that the average value of TDN content was 54.61 %. Maximum TDN content was observed in SSG 263 (55.25) followed by SSG 232 (55.09), SSG 256 (54.97) and SSG 241 (54.91) while, it was minimum in check SSG 59-3 (54.00). It means all the genotypes studied were found better than check for TDN content (Singh *et al.*, 2013).

**HCN content (ppm)**

As shown in Table-3, the difference among the genotypes with respect to the HCN content was significant at all the three cuts. The average values of HCN content were 279.06 at first cut, 233.85 at second cut and 173.13 at third cut. The desirable lower value was observed in check SSG 59-3 followed by SSG 222 and SSG 244 during all the three cuts. Maximum value exhibited by SSG 236 (318.33) at first cut, SSG 253 (278.33) at second cut and SSG 232 (193.33) at third cut (Singh *et al.*, 2013).

**Table 3 Mean values of green and dry fodder yield traits at different cut(s) in forage sorghum**

Genotype	Plant height (cm)			Leaves/plant			Stem girth (cm)			Tillers/plant		Fodder yield (g/plant/day)					
												Green			Dry		
	I Cut	II Cut	III Cut	I Cut	II Cut	III Cut	I Cut	II Cut	III Cut	I Cut	II Cut	I Cut	II Cut	III Cut	I Cut	II Cut	III Cut
SSG 222	240.3	123.6	63.3	12.8	9.0	6.6	4.5	3.1	2.1	1.3	2.3	2.2	1.2	0.4	0.6	0.5	0.2
SSG 224	240.3	176.3	103.3	10.5	9.1	6.3	3.6	3.4	2.3	2.0	3.2	2.9	1.3	0.8	0.8	0.5	0.4
SSG 225	255.0	155.6	103.3	11.9	9.5	6.6	4.0	3.4	2.4	1.8	2.7	2.3	1.4	0.7	0.7	0.6	0.3
SSG 226	239.6	167.6	115.0	11.5	10.3	8.0	3.8	3.5	2.7	2.4	3.6	3.3	1.5	0.9	1.0	0.6	0.4
SSG 227	244.0	155.3	71.6	11.3	9.3	7.3	3.8	3.6	2.9	1.3	1.9	3.3	1.1	0.5	0.8	0.4	0.2
SSG 231	232.3	149.6	58.3	10.9	9.2	7.0	4.6	4.0	2.4	1.5	2.6	3.9	1.2	0.5	1.0	0.5	0.2
SSG 232	243.3	160.0	88.3	11.5	10.2	7.3	4.0	3.9	2.9	1.3	2.5	3.0	1.0	0.4	0.8	0.4	0.2
SSG 233	256.0	101.6	47.6	11.2	8.3	6.6	4.2	3.6	2.7	1.5	2.1	2.2	1.1	0.4	0.6	0.3	0.2
SSG 234	245.3	134.3	88.0	10.8	9.0	5.6	4.0	3.5	2.0	1.5	2.8	3.0	1.4	0.3	1.0	0.4	0.1
SSG 236	238.0	135.6	88.3	10.9	8.6	6.6	4.2	3.5	2.4	1.4	2.4	2.8	1.1	0.4	0.8	0.4	0.2
SSG 241	271.3	141.3	94.3	11.7	9.2	7.3	4.2	3.7	2.6	1.4	2.5	2.2	1.3	0.9	0.6	0.4	0.3
SSG 244	228.3	130.6	94.0	12.2	9.9	7.6	4.0	3.3	3.0	1.2	1.5	3.5	1.6	0.5	0.9	0.5	0.2
SSG 253	253.6	139.6	113.3	11.5	9.3	7.3	4.4	3.6	3.4	1.8	3.1	3.5	1.0	0.8	1.1	0.4	0.3
SSG 256	220.0	135.0	73.3	10.9	9.4	6.6	3.7	3.3	2.1	1.4	2.3	3.1	1.6	0.3	0.9	0.5	0.1
SSG 263	256.0	90.6	53.3	12.1	8.8	5.6	4.2	3.8	2.1	1.3	1.8	3.8	1.4	0.3	1.2	0.4	0.1
SSG 59-3	253.6	145.3	116.6	12.1	10.0	8.3	4.3	3.3	2.6	2.5	3.8	2.5	1.4	0.8	0.7	0.5	0.3
CD (P=0.05)	34.8	54.2	30.8	1.0	1.3	1.7	0.5	0.6	0.6	0.3	0.7	1.3	0.6	0.3	0.4	0.8	0.8

**Table 4 Mean values of quality traits at different cut(s) in forage sorghum.**

Genotype	Nitrogen in plant (%)			Crude protein (%)	Crude fibre (%)	Ether extract (%)	Ash (%)	N free extract (%)	TDN (%)	HCN (ppm)		
	I Cut	II Cut	III Cut	I Cut	I Cut	I Cut	I Cut	I Cut	I Cut	I Cut	II Cut	III Cut
SSG 222	1.35	1.14	0.92	8.45	31.23	1.82	8.20	50.30	54.13	218.33	181.67	141.67
SSG 224	1.22	1.03	0.82	7.65	31.66	1.79	7.79	51.11	54.48	243.33	206.67	166.67
SSG 225	1.09	0.94	0.77	6.79	32.15	1.77	7.67	51.62	54.68	250.00	210.00	175.00
SSG 226	1.31	1.14	0.96	8.18	30.60	1.80	8.39	51.03	54.05	246.67	201.67	161.67
SSG 227	1.18	1.00	0.83	7.38	31.06	1.72	8.00	51.84	54.37	310.00	243.33	190.00
SSG 231	1.10	0.94	0.78	6.85	32.36	1.72	7.32	51.71	54.86	315.00	266.67	185.00
SSG 232	1.02	0.83	0.62	6.38	32.55	1.69	7.03	52.35	55.09	300.00	248.33	193.33
SSG 233	1.28	1.07	0.88	8.03	30.78	1.80	7.62	51.17	54.18	251.67	216.67	180.00
SSG 234	1.10	0.86	0.71	6.92	31.58	1.70	7.15	52.31	54.72	315.00	268.33	188.33
SSG 236	1.03	0.83	0.69	6.45	32.50	1.66	7.36	52.03	54.87	318.33	268.33	175.00
SSG 241	0.95	0.81	0.64	5.94	32.85	1.67	7.41	52.17	54.91	316.67	256.67	173.33
SSG 244	1.16	0.97	0.76	7.29	31.96	1.81	8.15	50.79	54.33	240.00	195.00	158.33
SSG 253	1.13	0.90	0.68	7.06	31.94	1.77	7.43	51.81	54.78	308.33	278.33	188.33
SSG 256	1.03	0.87	0.70	6.49	32.84	1.68	7.20	51.86	54.97	311.67	258.33	173.33
SSG 263	0.89	0.75	0.62	5.54	33.50	1.65	6.93	52.39	55.25	305.00	266.67	185.00
SSG 59-3	1.41	1.20	0.96	8.79	30.65	1.83	8.33	50.40	54.00	215.00	175.00	135.00
CD(P=0.05)	0.06	0.05	0.05	0.41	0.79	0.03	0.51	0.58	0.13	18.47	14.06	12.03

## CONCLUSION

Genetic variations in genotypes induced significant changes in morphological and yield traits. Data also suggested that new genotypes have potential to serve the forage purposes. Under the light of present study, the mutant genotypes SSG 226 is recommended for approval for general cultivation as it has better performance for fresh and dry matter yield. Furthermore, the variety SSG 226 appears to be leafier and therefore its dry

matter has the best nutritional value. Besides, two mutant genotypes SSG 231 and SSG 222 was also good for fodder yield and quality, respectively but it perform poorer for vice-versa. The future research should be to create more variability of check variety SSG 59-3 though inter-mating of these mutant lines specifically SSG 226, SSG 231 and SSG 222 to improve yield potential and nutritive value of forage sorghum.

## REFERENCES

- Al-Din A.A., Hessein E.M. and Ali E.A. 2012. Path coefficient assessment of yield and yield associated traits in sorghum (*Sorghum bicolor* L.) genotypes. *American-Eurasian Journal of Agricultural and Environmental Science*. **12** (6): 815-819.
- Alias, M.A., Bukhsh H.A., Ahmad R., Malik A.U., Hussain S. and Ishaque M. 2010. Agro-physiological traits of three maize hybrids as influenced by varying plant density. *Journal of Animal and Plant Sciences*. **20**(1): 34-39.
- AOAC, 1975. Official methods of analysis. 12<sup>th</sup> Ed. Association of Official Analytical Chemical, Washington, D. C.
- Ayub, M., Khalid M., Tariq M., Elahi M. and Nadeem M.A. 2012. Comparison of sorghum genotypes for forage production and quality. *Journal of Animal and Plant Sciences*. **22**(3): 733-737.
- Ayub, M., Nadeem M., Thir M., Ahmed Z. and Naem M. 2010. Evaluation of some forage sorghum varieties under the condition of central region, Saudi Arabia. *Annals Agricultural Sciences*. **39**(2): 649-654.

- Azam, M., Waraich E.A., Pervaiz A., and Nawaz F. 2010. Response of a newly developed fodder sorghum (*Sorghum bicolor* L. Monech) variety (F-9917) to NPK application. *Pakistan Journal of Life Sciences*. **8**(2): 117-120.
- Bibi A., Sadaqat H.A., Tahir M.H.N., Usman B.F. and Ali B. 2012. Genetic analysis of forage quality traits in sorghum-sudangrass hybrids under water stress. *Journal of Animal and Plant Sciences*. **22**(4): 1092-1100.
- Chughtai, S.R., Fateh J., Munawwar M.H., and Hussain M. 2007. Participatory sorghum varietal evaluation, selection in Pakistan. *Pakistan Journal of Agricultural Research* **55**: 19-26.
- Devendra, C., and Leng R.A. 2011. Feed resources for animals in Asia: issues, strategies for use, intensification and integration for increased productivity. *Asian-Australian Journal of Animal Sciences*. **24** (3): 303 – 321.
- Ghasemi A., Karim M.H. and Ghasemi M.M. 2012. Green fodder yield performance of different varieties of sorghum grown in an arid region. *International Journal of Agriculture and Crop Sciences*. **4**(13): 839-843.
- Hogg P.G. and Ahlgren H.L. 1942. A rapid method for determining HCN content of single plant of sudan grass. *Journal of American Society of Agronomy*. **43**: 199-200.
- Hussain, A., Muhammad D., Khan S., and Bhatti M.B. 1995. Performance of various cultivars of forage sorghum under rainfed conditions. *Journal of Agricultural Research*. **33**: 413-418.
- Mukherjee A.K., Malti S. and Nanda M.R. 1998. Strategy for maximization of forage production. *National Seminar on Strategy for maximization of forage production by 2000 A.D. held at BCKV, West Bengal*. 5-7.
- Nabi, C.G., Riaz M. and Ahmed G. 2006. Comparison of some advanced lines of *Sorghum bicolor* for green fodder/dry matter yield and morpho economical parameters. *Journal of Agricultural Research*. **44**: 191-196.
- Seetharam, K. and Ganeshamurthy, K. 2013. Characterization of sorghum genotypes for yield and other agronomic traits through genetic variability and diversity analysis. *Electronic Journal of Plant Breeding*. **4**(1): 1073-1079.
- Singh J., Ranwah B.R., Chaudhary L., Lal C., Dagla, M.C. and Kumar, V. 2013. Evaluation for genetic variability, correlation and path coefficient in mutant population of forage sorghum (*Sorghum bicolor*). *The Bioscan*. **8** (4): 1471-1476.
- Steel, R.G.D., Torrie J. H. and Dickey D.A. 1997. Principles and Procedures of Statistics. A biometrical approach. 3rd Ed. McGraw Hill, Inc. Book Co. N.Y. (U.S.A.): 352-358.
- SIA. 2013. State of Indian Agriculture report.pp23-29
- Tauqir, N.A., Sarwar M., Jabbar M.A., and Mahmood S. 2009. Nutritive value of jumbo grass (*Sorghum bicolor sorghum sudanefe*) silage in lactating niliravi buffaloes. *Pakistan Veterinary Journal*. **29**: 5-10.
- Ullah, A., Khan A.A., Nawab K., Khan A. and Islam B. 2007. Growth characters and fodder production potential of sorghum varieties under irrigated conditions. *Sarhad Journal of Agriculture*. **23**(2): 265- 268.
- Yousef, E., Carmi, A., Nikbachat, M., Zenou, A., Umiel, N. and Miron, J. 2009. Characteristics of tall versus short-type varieties of forage sorghum grown under two irrigation levels, for summer and subsequent fall harvests, and digestibility by sheep of their silages. *Animal Feed Science Technology*. **152**: 1-11.